## CHILDREN'S Diphenhydramine (BENADRYL) (12.5mg/5ml) Do not exceed 4 doses in 24 hours

Weight	Dose	Dosing Interval
20 pounds	<sup>3</sup> ⁄ <sub>4</sub> - 1 teaspoon	Every six hours
25 pounds	1 teaspoon	Every six hours
30 pounds	1 – 1.5 teaspoons	Every six hours
35 pounds	1.5 – 1.75 teaspoons	Every six hours
40 pounds	1.75 teaspoons	Every six hours
45 pounds	2 teaspoons	Every six hours

Acetaminophen (tylenol)
Dosage interval is every four hours (Do not exceed 5 doses in 24 hours)
Do not use before 2 months of age. Between 2 and 3 months of age, please consult a doctor

						, I			
Age	0-3 mos	4-11 mos	12-23 mos	2-3 yrs	4-5 yrs	6-8 yrs	9-10 yrs	11 yrs	12 yrs
Weight	6-11 lbs	12-17 lbs	18-23 lbs	24-35 lbs	36-47 lbs	48-59 lbs	60-71 lbs	72-95 lbs	95lbs+
Children & Infant Acetominoph en Suspension (160mg/5ml)	1.25ml (CALL doctor FIRST)	2.5 ml or 1/2 tsp	3.75 ml or 3/4 tsp	1 tsp or 5ml	1.5 tsp or 7.5ml	2 tsp or 10ml	2.5 tsp or 12.5ml	3 tsp	
Children's Meltaway Tablets (80mg each)				2 tablets	3 tablets	4 tablets	5 tablets	6 tablets	
Junior Acetominoph en Meltaway Tablets (160 mg)						2 tablets	2.5 tablets	3 tablets	4 tablets

Ibuprofen (motrin and advil)

Dosing interval is every 6 hours (do not exceed 4 doses in 24 hours)
Do not use before age 6 months

Age	0-5 mos	6-11 mos	12-23 mos	2-3 yrs	4-5 yrs	8-6 yrs	9-10 yrs	11 yrs
Weight	6-11 lbs	12-17 lbs	18-23 lbs	24-35 lbs	36-47 lbs	48-59 lbs	60-71 lbs	72-95 lbs
Concentrat ed Ibuprofen Infants' Drops (50mg/ 1.25ml)		1.25 ml	1.875 ml					

Children's ibuprofen Suspensio n (100mg/ 5ml)	 	 1 tsp	1.5 tsp or 7.5ml	2 tsp or 10 ml	2.5 tsp or 12.5ml	3 tsp or 15ml
Motrin Junior Strength Easy-to- Swallow Caplets (100mg)	 	 		2 caplets	2.5 caplets	3 caplets
Motrin Junior Strength Chewable Caplets (100mg)	 	 1 tablet	1.5 tablets	2 tablets	2.5 tablets	3 tablets