

CHILDREN'S Diphenhydramine (BENADRYL) (12.5mg/5ml)

Dosage interval is every six hours. Do not exceed 4 doses in 24 hours

Child's weight (pounds)	20-24 lbs	25-37 lbs	38-49 lbs	50-99 lbs	100 + lbs	
Liquid 12.5 mg/5 ml	4	5	7.5	10	20	ml
Liquid 12.5 mg/1 tsp	$\frac{3}{4}$	1	1 $\frac{1}{2}$	2	4	tsp
Chewable 12.5 mg	--	1	1 $\frac{1}{2}$	2	4	tablets
Tablets 25 mg	--	$\frac{1}{2}$	$\frac{1}{2}$	1	2	tablets
Capsules 25 mg	--	--	--	1	2	caps

Acetaminophen (tylenol)

Dosage interval is every four hours (Do not exceed 5 doses in 24 hours)

Do not use before 2 months of age. Between 2 and 3 months of age, please consult a doctor

Age	0-3 mos	4-11 mos	12-23 mos	2-3 yrs	4-5 yrs	6-8 yrs	9-10 yrs	11 yrs	12 yrs
Weight	6-11 lbs	12-17 lbs	18-23 lbs	24-35 lbs	36-47 lbs	48-59 lbs	60-71 lbs	72-95 lbs	95lbs+
Children & Infant Acetaminophen Suspension (160mg/5ml)	1.25ml (CALL doctor FIRST)	2.5 ml or 1/2 tsp	3.75 ml or 3/4 tsp	1 tsp or 5ml	1.5 tsp or 7.5ml	2 tsp or 10ml	2.5 tsp or 12.5ml	3 tsp	---
Children's Meltaway Tablets (80mg each)	--	--	---	2 tablets	3 tablets	4 tablets	5 tablets	6 tablets	---
Junior Acetaminophen Meltaway Tablets (160 mg)	---	---	---	---	---	2 tablets	2.5 tablets	3 tablets	4 tablets

Ibuprofen (motrin and advil)

Dosing interval is every 6 hours (do not exceed 4 doses in 24 hours)

Do not use before age 6 months

Age	0-5 mos	6-11 mos	12-23 mos	2-3 yrs	4-5 yrs	8-6 yrs	9-10 yrs	11 yrs
Weight	6-11 lbs	12-17 lbs	18-23 lbs	24-35 lbs	36-47 lbs	48-59 lbs	60-71 lbs	72-95 lbs
Concentrated Ibuprofen Infants' Drops (50mg/ 1.25ml)	---	1.25 ml	1.875 ml	---	---	---	---	---

Children's ibuprofen Suspension (100mg/ 5ml)	---	---	---	1 tsp	1.5 tsp or 7.5ml	2 tsp or 10 ml	2.5 tsp or 12.5ml	3 tsp or 15ml
Motrin Junior Strength Easy-to- Swallow Caplets (100mg)	---	---	---	---	---	2 caplets	2.5 caplets	3 caplets
Motrin Junior Strength Chewable Caplets (100mg)	---	---	---	1 tablet	1.5 tablets	2 tablets	2.5 tablets	3 tablets