

## Acetaminophen (Tylenol®) Dosage

Dosage interval is every 4 hours. Do not exceed 5 doses in 24 hours.

Do not use before 2 months of age. Between 2 and 3 months of age, please consult a doctor.

Age	Weight	Infant & Children's Suspension Liquid (160mg/5ml)	Children's Soft Chew Tablets (80mg each)	Junior Strength Chewable Tablets (160mg each)
0-3 mos	6-11 lbs	1.25ml (CALL doctor FIRST)	--	--
4-11 mos	12-17 lbs	2.5ml or ½ tsp	--	--
12-23 mos	18-23 lbs	3.75ml or ¾ tsp	--	--
2-3 yrs	24-35 lbs	5ml or 1 tsp	2 tablets	--
4-5 yrs	36-47 lbs	7.5ml or 1.5 tsp	3 tablets	--
6-8 yrs	48-59 lbs	10ml or 2 tsp	4 tablets	2 tablets
9-10 yrs	60-71 lbs	12.5ml or 2.5 tsp	5 tablets	2.5 tablets
11 yrs	72-95 lbs	15ml or 3 tsp	6 tablets	3 tablets
12 yrs	96+ lbs	--	--	4 tablets

## Ibuprofen (Motrin® and Advil®) Dosage

Dosage interval is every 6 hours. Do not exceed 4 doses in 24 hours.

Do not use before 6 months of age.

Age	Weight	Concentrated Infants' Drops (50mg/1.25ml)	Children's Ibuprofen Suspension (100mg/1tsp)	Junior Strength Chewables (100mg each)
6-11 mos	12-17 lbs	1.25ml	--	--
12-23 mos	18-23 lbs	1.875ml	--	--
2-3 yrs	24-35 lbs	--	5 ml or 1 tsp	1 tablet
4-5 yrs	36-47 lbs	--	7.5ml or 1.5 tsp	1.5 tablets
6-8 yrs	48-59 lbs	--	10ml or 2 tsp	2 tablets
9-10 yrs	60-71 lbs	--	12.5ml or 2.5 tsp	2.5 tablets
11 yrs	72-95 lbs	--	15ml or 3 tsp	3 tablets

## Diphenhydramine (Benadryl®) Dosage

Dosage interval is every 6 hours. Do not exceed 4 doses in 24 hours.

Weight	Dose
20-24 lbs	¼ - 1 tsp
25-29 lbs	1 tsp
30-34 lbs	1 – 1.5 tsp
35-39 lbs	1.5 – 1.75 tsp
40-44 lbs	1.75 tsp
45 lbs	2 tsp